


Boldenone

Boldenone Undecylenate Injection, USP

DIN 00089384

200 mg/mL

10 mL Multiple-Dose vial, Cartons of 1 vial



Systematic name: (8R,9S,10R,13S,14S,17S)-17-hydroxy-10,13-dimethyl-6,7,8,9,10,11, 12,13,14,15,16,17-dodecahydro-3H-cyclopenta[a]phenanthren-3-one

DESCRIPTION

Boldenone Undecylenate is a 17 β -hydroxy-androsta-1,4-dien- 3-one-10- undecenoate steroid ester which possess marked anabolic properties and a small amount of androgenic activity. Anabolic and androgenic agents became widely used in the treatment of certain pathophysiological or catabolic processes. In many instances, it is desirable to maintain a constant level of effect over a long period of time. Boldenone Undecylenate is a long-acting injectable agent which has a rapid onset of action.

ATHLETISME

Boldenone Undecylenate is a very popular steroid. Boldenone is a highly anabolic, yet moderately androgenic steroid. For this very reason, it is typically taken with other steroids like testosterone during a mass cycle. The main benefit of taking Boldenone Undecylenate is that it increases protein synthesis of the muscle cells. Boldenone gives you slower muscles gains but of much higher quality as opposed to the normal quick muscle gains from the testosterone. It is suggested to ask for professional advices before using this steroid. We expect a weight gain of 20 lbs in 6 weeks. In around 3 weeks, results could be seen. These are not staggering, but the gains will be more permanent than any ones provided by other available testosterones. This steroid stays active in the system longer than most of the testosterones as well. This makes Boldenone a poor choice if there is a possibility of being drug tested. Boldenone is very common in the pre-contest arena for two main reasons. It contains a low amount of aromatization and the water retention caused by the product is insignificant. This makes Boldenone a good pre-contest steroid. Boldenone is well known to give a good increase in the pumps while working out. This is due to the increase in red blood cells provided by this steroid. It is also well known for causing a dramatic increase in appetite. When taken with a good mass building steroid, this is a guaranteed formula for successful gains in muscle mass.

COMPOSITION Medicinal ingredients and Non-Medicinal ingredients.

Each mL contains: Boldenone undecylenate, 200 mg; benzyl alcohol, 9 mg; benzyl benzoate, 0.1 mL; Cottonseed oil USP. q.s

CLINICAL PHARMACOLOGY ACTIONS

Studies conducted in laboratory animals to evaluate the pharmacological activity characterized Boldenone Undecylenate as having distinct anabolic properties together with a certain degree of androgenic activity. It does not have marked antigonadotrophic properties, nor does it produce any clear-cut effects on the endometrium, conditions that are commonly observed when similar substances are used. In clinical trials at the recommended dosage, Boldenone Undecylenate had a marked anabolic effect in debilitated users. Appetite improvement, weight gain, increased vigor and improvement was noted in musculature. This would be expected with an anabolic agent such as Boldenone Undecylenate, particularly where there had been marked tissue breakdown associated with disease, prolonged anorexia or overwork.

INDICATIONS

Boldenone Undecylenate is recommended as an aid for treating debilitated users when an improvement in weight or general physical condition is desired. Debilitation often follows disease or may occur following overwork and over-exertion. Boldenone Undecylenate improves the general state of debilitated users, thus aiding in correcting weight losses and improving appetite. It is not a substitute for a well-balanced diet. Optimal results can be expected only when good management and feeding practices are utilized. Boldenone Undecylenate should be considered only as adjunctive therapy to other specific and supportive therapy for diseases, surgical cases, and traumatic injuries.

CONTRAINDICATIONS

Androgenic steroids are contraindicated in men with carcinomas of the breast or with known or suspected carcinomas of the prostate and in women who are or may become pregnant. Higher doses of Boldenone Undecylenate Injection can cause virilization symptoms such as deep voice, increased production of the sebaceous gland and acne. Boldenone Undecylenate Injection can increase hair growth on the face and legs. Boldenone Undecylenate is contraindicated if you have a history of hypersensitivity to any of its components.

ADVERSE REACTIONS

Treatment with Equipoise (Boldenone Undecylenate injection) may result in androgenic effects (masculinization, behavioural effects) being expressed in some users. If these effects occur, they may persist for up to 8 to 10 weeks.

DOSAGE AND ADMINISTRATION

Boldenone Undecylenate should be administered every 7 to 10 days at a dose of 400 mg to 600 mg for 10 weeks of therapy. Overdosage An overdose of Boldenone Undecylenate injection may result in prolonged androgenic effects.

STORAGE INSTRUCTIONS

Vial should be kept away from light and stored in controlled temperature from 20-25 degree Celsius (68° TO 77°F). Warming and shaking the vial should redissolve any crystals that may have formed during storage. Keep out of reach of children.

